



Blake, Holly and Alsaleh, Eman and Windle, Richard (2014) Behavioural intervention to increase physical activity among patients with coronary heart disease (CHD): randomised controlled trial. In: UK Society for Behavioural Medicine 10th Annual Scientific Meeting, 3-4 December 2014, Nottingham, UK.

Access from the University of Nottingham repository:

<http://eprints.nottingham.ac.uk/39016/1/2014%20Blake%20et%20al%20Cardiac%20Jordan%20UKSBM.pdf>

Copyright and reuse:

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the University of Nottingham End User licence and may be reused according to the conditions of the licence. For more details see:
http://eprints.nottingham.ac.uk/end_user_agreement.pdf

A note on versions:

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact eprints@nottingham.ac.uk

Citation: BLAKE, H, ALSALEH, E and WINDLE, R, 2014. Behavioural intervention to increase physical activity among patients with coronary heart disease (CHD): randomised controlled trial. In: Proceedings of the 10th UK Society for Behavioural Medicine Annual Scientific Meeting, 3-4 December, Nottingham.

Behavioural intervention to increase physical activity among patients with coronary heart disease (CHD): randomised controlled trial Holly Blake, Eman Alsaleh, Richard Windle

University of Nottingham, Nottingham, UK

Background: Patients with CHD often do not follow prescribed physical activity recommendations.

Aim: To assess the efficacy of behavioural intervention to increase physical activity in patients not attending structured cardiac rehabilitation programmes.

Design: Randomised controlled trial comparing 6 month multi-component behavioural change intervention (n=85) with usual care (n=71). Intervention included face-to-face individualised consultation and telephone support (for goal-setting, feedback and self-monitoring) and reminder text messages.

Setting: Two hospitals in Jordan, Middle East.

Participants: 156 patients with CHD (mean age 57.5 years; 54% male, 46% female).

Measurements: Outcomes measured at baseline and 6 months. Primary outcome was physical activity. Secondary outcomes were blood pressure, body mass index, exercise self-efficacy and health-related quality of life.

Findings: Intervention and control groups were comparable at baseline. Moderate PA significantly increased in intervention group compared with control group (mean change (SD) of frequency: 0.23 (0.87) days/week versus -.06 (0.40), duration: 15.53 (90.15) minutes/week versus -3.67 (22.60) minutes/week and intensity: 31.05 (105.98) Metabolic equivalents (METs) versus 14.68 (90.40) METs. Walking significantly increased in the intervention group compared with control group (the mean change (SD) of frequency: 3.15 (2.75) days/week versus 0.37 (1.83) days/week, duration: 150.90 (124.47) minutes/week versus 24.05 (195.93) minutes/week and

intensity: 495.12 (413.74) METs versus 14.62 (265.06) METs. Intervention participants had significantly lower blood pressure, lower body mass index, greater exercise self-efficacy and better health-related quality of life compared with controls.

Conclusions: Multi-component behavioural intervention increases physical activity, improves body composition and improves psychological outcomes in CHD patients not attending structured rehabilitation programmes.